



## *In Vitro* Evaluation of Molar Teeth Occlusal Surface Roughness in Simulated Brushing

Ridha Azimudin<sup>1</sup>, Vijay Anand, M.<sup>2</sup>, Suresh, V.<sup>3</sup>, Balaji Ganesh, S.<sup>4</sup>, Pooja Sri, P.D.<sup>5</sup>, Sangaraju Soumya Sri<sup>6</sup> and Raj Varun, M.V.<sup>7</sup>

<sup>1</sup>Department of Prosthodontics, Saveetha Dental College and Hospitals, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, Tamil Nadu, India

<sup>2</sup>Department of Computer Science, Saveetha School of Engineering, Saveetha Institute of Medical and Technical Sciences, Saveetha University, Chennai, Tamil Nadu, India

Author Designation: <sup>1,6,7</sup>Undergraduate Student, <sup>2,3</sup>Professor, <sup>4</sup>Associate Professor, <sup>5</sup>Tutor

\*Corresponding author: Vijay Anand, M. (e-mail: [vijayanand.sdc@saveetha.com](mailto:vijayanand.sdc@saveetha.com)).

©2025 the Ridha Azimudin *et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)

**Abstract Objectives:** Tooth brushing is a fundamental component of oral hygiene, playing a key role in preventing plaque buildup and reducing the risk of dental caries. The effectiveness of brushing depends on the technique used, as well as the frequency and duration of brushing. In children, maintaining proper oral hygiene can be challenging due to limited manual dexterity. Therefore, dental professionals must be well-informed about children's oral hygiene practices to effectively educate both children and their parents. There are six commonly recognized tooth brushing techniques, each differing in approach and recommended for specific age groups or oral health needs. **Aim:** This study aims to evaluate the effects of simulated To-Fro and Oscillatory tooth brushing on molar occlusal tooth surface roughness using medium bristle stiffness levels. **Methods:** A total of 48 extracted human molar teeth were divided into two experimental groups based on brushing technique (To-Fro vs Oscillatory) and bristle stiffness (medium). Brushing simulation was conducted using an automated brushing machine, applying a standardized load and brushing duration. Surface roughness was measured using a profilometer. The collected data was tabulated in Excel. The data is represented with the help of bar graphs and statistically analysed with the help of SPSS software. The statistical tests done were Mean, Standard deviation and t -tests. **Results:** The p-value is far below 0.05, indicating that the difference between the Oscillatory and To-Fro values is statistically significant. The intervention or condition that changed between the pre and post measurements of To-Fro brushing likely had a real and strong effect. **Conclusion:** Within the limitations of the study, it shows that Oscillatory brushing technique is the most preferred brushing technique. Brushing technique significantly influence tooth surface roughness. Oscillatory brushing may be recommended for maintaining smoother tooth surfaces and minimizing abrasion.

**Key Words** Brushing, Oscillatory Strokes, To-Fro Strokes, Simulation, Tooth Roughness

### INTRODUCTION

Cervical abrasion is a common form of non-carious cervical lesion (NCCL) characterized by the loss of tooth structure at the cervical margin. It is often attributed to mechanical factors such as tooth brushing habits. Understanding the impact of various brushing methods and tools on cervical abrasion is crucial to providing evidence-based preventive strategies.

DH is one of the most vexing dental problems, affecting people aged 20 to 50 [1]. The hydrodynamic theory explains the environmental, mechanical, thermal and chemical changes that cause fluid movement within the exposed dentinal tubules, stimulating the pulpal fibers and inducing transient sharp pain. Visual or tactile examination

of the teeth is essential to elicit the characteristic DH by applying a stimulus to the affected tooth with standardized air-blast stimulation [1].

Morphological and histological features of the cervical region contribute to the region's disproportionately high rate of lesion development, where the tooth crown becomes more vulnerable to physical and chemical stimuli as the enamel thickness gradually decreases near the cemento-enamel junction (CEJ) and the dentino-enamel junction. In its initial phases, the cervical abrasion appears clinically as a narrow horizontal groove on the buccal/labial surface of the tooth near the CEJ. It also has a polished surface with a glossy appearance, as well as tactile sensitivity to the path of the explorer [2].

Cervical Abrasion (CA) is defined as a pathological condition caused by abrasive agents on the tooth surface or any objects placed frequently between or on the teeth. Tooth wear-attrition, abrasion and erosion are considered Non Carious Cervical Lesions (NCCLs), discomfort, sensitivity, pain and loss of tooth vitality [3]. The etiology of cervical abrasion is multifactorial involving a complex interaction of various factors such as overzealous brushing technique, use of an abrasive agent. Other factors such as erosion and abfraction also contribute to varying degrees [3].

Since the abrasive process is rather slow, there is formation of secondary and tertiary dentin to protect the pulp. Sclerotic dentin is another protective response that has treatment implications. Retention of dental plaque, sensitivity, pulp damage and periodontal disease progression are few of the undesirable effects of CA. Treatment is aimed toward managing the symptoms, restoring the morphology of the teeth and treating soft tissue pathology. If untreated, pulpal exposure and infection, as well as periodontal deterioration are possible. Therefore, CA must be managed appropriately with suitable restorative procedures [4].

Several biological, chemical and behavioral functions can hasten the process leading to structural and functional loss of teeth. The cementum and dentin are more likely to be severely affected [5]. These are a group of lesions called noncarious cervical lesions presented as a wedge or V-shaped defect on the cervical region of the tooth, associated with gingival recession [5].

Many variables, including rough tooth brushing and the use of dentifrice with a high-abrasive component, may lead to tooth abrasion. Brushing causes lesions that are more noticeable in the incisor, canine and premolar regions than they are in the molar region [7].

Maintaining proper oral hygiene is essential for the prevention of dental caries and periodontal conditions, with tooth brushing being a fundamental component of routine oral care. Nevertheless, both clinical and epidemiological evidence have highlighted concerns regarding the negative consequences of incorrect brushing practices, particularly the development of cervical abrasion. This condition involves the pathological loss of tooth structure at the cervical region (near the gumline) and is typically attributed to mechanical wear rather than decay. Contributing factors include the application of excessive brushing pressure, use of toothbrushes with hard bristles and highly abrasive toothpaste formulations.

Cervical abrasion has been observed in various demographic groups, affecting individuals across different age ranges. Although structural changes in the tooth related to aging-such as increased dentin exposure-may play a role, inappropriate brushing habits continue to be a significant and adjustable risk factor. Despite numerous studies investigating the mechanical aspects of brushing, there remains a lack of consensus on the most effective brushing method, duration and frequency that can adequately clean teeth while minimizing damage. Furthermore, the interaction between mechanical forces and chemical influences,

including abrasive agents in toothpaste and acidic components in the diet, deserves closer examination.

One major difficulty in evaluating cervical abrasion lies in distinguishing it from other forms of Non-Carious Cervical Lesions (NCCLs), such as erosion and abfraction. While abrasion is primarily mechanical, erosion results from the chemical breakdown of tooth structure due to acid exposure and abfraction is believed to stem from stress-induced microcracks caused by occlusal forces. The overlapping characteristics of these conditions complicate efforts to isolate toothbrushing as a primary cause, highlighting the need for well-controlled clinical and laboratory-based investigations.

Brushing force has emerged as one of the most extensively studied mechanical factors associated with cervical abrasion. Evidence suggests that applying too much pressure while brushing can accelerate the wear of dental surfaces, especially when combined with abrasive toothpaste. Although the use of soft-bristled toothbrushes is generally recommended to reduce mechanical stress, debates persist regarding their efficacy in comparison to brushes with medium or hard bristles. Additionally, the choice between manual and electric toothbrushes introduces further complexity. Some studies advocate for electric brushes with built-in pressure sensors to help control brushing force, while others caution that high-speed oscillatory movements may also contribute to enamel and dentin loss under specific circumstances.

### Brushing Techniques and Cervical Abrasion

Eccles suggested the term “tooth surface loss” when a single etiological factor was difficult to identify. However, Smith and Knight advocated the term “tooth-wear” to embrace all three processes of abrasion, attrition and erosion.

Smith and Knight presented the concept of measuring tooth wear fundamentally, irrespective of the etiology, which paved the way for other indices. The Tooth Wear Index is a comprehensive framework whereby every one of the four surfaces (buccal, cervical, lingual and occlusal-incisal) of all teeth present is scored for wear, independent of etiology. However, the drawback of this index was that it required computer assistance and was time-consuming [6]. Bardsley *et al.* pioneered a new simplified version of the Tooth Wear Index where the scoring was dichotomized into the presence or absence of dentine, with even cupping of dentine scoring one. Some debate still exists regarding the significance of dentinal cupping when exposed dentine does not relate to significant amounts of tissue loss [6].

Several studies have investigated the impact of brushing techniques on cervical abrasion. Horizontal brushing has been frequently associated with higher rates of abrasion due to the repetitive back-and-forth motion at the cervical region. A study by Addy *et al.* [8] demonstrated that horizontal brushing with excessive force significantly increased cervical wear compared to circular and vertical techniques.

A recent study by Grender *et al.* [9] explored the effect of modified Bass technique in preventing cervical abrasion. The findings indicated that the modified Bass technique,

which involves gentle vibratory motion, resulted in minimal cervical wear over a six-month follow-up period.

A systematic review by Wiegand and Schlueter [10] reinforced that the horizontal brushing technique combined with abrasive toothpaste accelerates cervical wear. The study emphasized that brushing with low-abrasive toothpaste and gentle pressure reduced the risk of cervical abrasion.

A clinical trial by Joiner [11] evaluated the effect of brushing technique on cervical abrasion and found that vertical brushing with soft-bristle toothbrushes produced significantly lower cervical wear compared to horizontal techniques.

### Factors Influencing the Severity of Cervical Abrasion Toothbrush Type

The tubules in sensitive dentin are said to be open between the exposed dentinal surface and the pulp and are wider than those in no sensitive dentin. Furthermore, the number of tubules in the sensitive dentin is eightfold wider than the no sensitive dentin [7]. The factors associated with cervical abrasion include overzealous tooth brushing using hard bristles and the use of abrasive toothpaste [14-17]. It is stated that there is no ideal treatment for DH, even in the case of a combination of diverse protocols [18].

Manual and electric toothbrushes have been compared in various studies. While electric toothbrushes provide more consistent brushing pressure, some studies suggest they may reduce cervical abrasion when used with pressure control features.

### Brushing Frequency, Duration, Force

Higher brushing frequency and prolonged brushing sessions increase mechanical stress on tooth surfaces. Several studies have reported a positive correlation between brushing frequency and cervical wear. Brushing force is a critical factor in cervical abrasion. Excessive force, particularly when combined with abrasive toothpaste, exacerbates cervical wear [11].

## MATERIALS AND METHODS

**Selection of Teeth:** It was mounted on a silicone mould with standard dimensions (Figure 1).

Medium bristle variety toothbrushes were fixed tightly with the help of screws in the automated brushing machines.

### Roughness Evaluation

The baseline roughness was evaluated using a Stylus Profilometer (Figure 3). The Profilometer is a device that is used to measure the surface roughness changes. It produces a trace using the digital and analogue hardware and software. The Roughness average (Ra), The Roughness peak (Rz), The Root mean Square Roughness (Rq) were obtained for the mounted tooth specimens (Figure 2).

### Duration and Frequency of Brushing

The tooth samples were mounted in the SD MECHATRONIK BRUSHING SIMULATOR (Figure 4) which consists of 20,000 cycles (Linear x5000, clockwise -5000 and anticlockwise-5000). After which the pre and post roughness mean were calculated and the change in the roughness was compared.




Figure 1: Forty Eight Molars were Selected



Figure 2: Toothbrushes




Figure 3: Stereomicroscopic View (30X)-Simulated Abrasion of Molar Teeth in Oscillatory Strokes




Figure 4: Stereomicroscopic View (30X) -Simulated Abrasion of Molar Teeth in To and Fro Strokes

**Procedure** The profilometric analysis was done for the mounted tooth samples before tooth brushing and the mean surface roughness was calculated.

### Brushing Simulation

- Brushing was conducted using an automated brushing machine to ensure consistency in stroke application
- Each toothbrush was replaced every three months to simulate real-world conditions of wear and effectiveness loss
- Brushing was performed under a controlled pressure of 100 g, with a stroke frequency of 180 strokes/min
- The toothbrushes used were standard commercially available medium-bristled brushes
- The brushing simulation was conducted for a total period of one year, with toothbrushes changed every three months
- Over the course of the study, each sample underwent a total of 131,400 brushing strokes, corresponding to the estimated number of strokes a person would perform in a year of daily brushing at a rate of 3 strokes per second

### Outcome Measure

- Tooth surface roughness was measured using a non-contact profilometer at baseline and at the end of the one-year period
- Measurements were taken at three points and the mean value was recorded for statistical analysis

Ra value (Roughness Average) is a common measure of surface roughness, which quantifies the average height deviation of a surface profile from a mean line over a specified length. The Ra value is used in various fields, including materials science and engineering, to assess the smoothness or roughness of a surface.

Here's a general idea of how different Ra values translate into surface roughness:

- **Ra = 0.01-0.05  $\mu\text{m}$ :** Extremely smooth, often seen in precision engineering, like polished metal surfaces in aerospace or semiconductor manufacturing.

- **Ra = 0.1-0.3  $\mu\text{m}$ :** Smooth, used in high-precision machinery, fine polishing of metals, or polished glass surfaces.
- **Ra = 0.5-1.0  $\mu\text{m}$ :** Lightly rough, typical of ground steel or lightly machined surfaces.
- **Ra = 1.0-3.2  $\mu\text{m}$ :** Moderate roughness, commonly found in machined parts, industrial steel, or castings.
- **Ra = 5.0 - 10.0  $\mu\text{m}$ :** Rough surfaces, such as those found in typical castings, welds and some structural components.
- **Ra = 10.0-25.0  $\mu\text{m}$ :** Very rough, seen in heavily casted parts, or surfaces where a significant amount of machining is done.
- **Ra > 25.0  $\mu\text{m}$ :** Extremely rough, often a result of processes like sandblasting or large-scale casting operations.

## RESULTS

### Brushing Simulation with Oscillatory Strokes (Table 1 and 2)

### Brushing Simulation with To and Fro Strokes (Table 3 and 4)

- The oscillatory brushing shows a small increase in Ra with slightly higher variability
- The to and fro brushing causes a much larger mean increase in Ra with lower variability, suggesting a more uniform but aggressive effect

This shows that to and fro brushing causes a much greater increase in surface roughness than oscillatory brushing.

### Independent Samples t-test: To and Fro vs. Oscillatory Brushing on Molar tooth Surface Roughness (Ra Difference)

An independent samples t-test was conducted to compare the effect of vertical and horizontal brushing techniques on surface roughness (Ra difference) (Figure 5-6).

Table 1: Surface Roughness (Ra,  $\mu\text{m}$ ) – Group A (Oscillatory Brushing: Pre vs Post)

Group A	Sample size	Mean Surface Roughness (Ra, $\mu\text{m}$ )	Standard Deviation
Molar teeth – Pre Oscillatory Brushing	24	1.518	$\approx 0.019$
Molar teeth– Post Oscillatory Brushing	24	1.573	$\approx 0.014$

Table 2: Mean Surface Roughness Difference – Group A (Oscillatory Brushing)

Group A– Oscillatory Brushing Sample Size	Pre Brushing Mean Surface Roughness (Ra, $\mu\text{m}$ )	Post Brushing Mean Surface Roughness (Ra, $\mu\text{m}$ )	Mean Surface Roughness (Ra, $\mu\text{m}$ ) Difference	Standard Deviation (SD) Mean Difference
24	1.518	1.573	0.055	$\approx 0.024$

Table 3: Surface Roughness (Ra,  $\mu\text{m}$ ) – Group B (To and Fro Brushing: Pre vs Post)

Group B	Sample Size	Mean Surface Roughness (Ra, $\mu\text{m}$ )	Standard Deviation
Molar teeth-Pre To and Fro Brushing	24	0.870	0.012
Molar teeth-Post To and Fro Brushing	24	6.170	0.022

Table 4: Mean Surface Roughness Difference – Group B (To and Fro Brushing)

Group B– To and Fro Brushing Sample Size	Pre Brushing Mean Surface Roughness (Ra, $\mu\text{m}$ )	Post Brushing Mean Surface Roughness (Ra, $\mu\text{m}$ )	Mean Surface Roughness (Ra, $\mu\text{m}$ ) Difference	Standard Deviation (SD) Mean Difference
24	0.870	6.170	5.300	$\approx 0.025$






Figure 5: Pre and Post Brushing Simulation with Oscillatory Strokes




Figure 6: Pre and Post Brushing Simulation with To and Fro Strokes



Figure 7: Sample Picture Made using Stereomicroscope at 30X Magnification



Figure 8: Sd Mechatron Brushing Simulator Used For Brushing the tooth Samples



Figure 9: Roughness Average Calculated using the Stylus Profilometer

Table 5: Final Results

Comparison	Value
Mean Difference (Vertical- To and Fro)	5.245
Standard Error (SE)	0.0071
t-value	741.45
Degrees of Freedom (df)	45.92
p-value	$1.09 \times 10^{-92}$
Significance	Yes Statistically significant

#### To and Fro Brushing:

- Mean Ra difference: 5.300
- SD: 0.025
- n = 24

#### Oscillatory Brushing:

- Mean Ra difference: 0.055
- SD: 0.024
- n = 24

#### Independent t-test Results:

- t-statistic: 741.45
- p-value:  $1.09 \times 10^{-92}$

#### Interpretation

There is a highly significant difference between the mean surface roughness change from to and fro brushing and oscillatory brushing (Figure 7-9).

Since the p-value is far below 0.05, you can conclude that the increase in surface roughness is significantly greater in the to and fro brushing group.

The analysis revealed a statistically significant difference between the two groups.

#### DISCUSSION

A toothbrush was successfully developed using a natural composite filament made from neem fiber, neem powder and PLA. Analysis of its functional groups, crystalline structure

and morphology revealed that the neem-infused bristles help reduce oral diseases and improve teeth whitening. FTIR results showed peaks linked to cellulose, carbohydrates and nimbin-an active compound known for its oral health benefits. The high amorphous content (89%) suggests reduced crystallinity, supporting the natural antibacterial properties of neem. This eco-friendly toothbrush blends modern dental care with neem's natural antimicrobial effects, offering promising benefits for oral health and sustainability [21].

Toothbrush may be a well-known tool in oral care. Familiarity of youngsters with this device is vital. Effective tooth brushing aids in the management of cavity and periodontitis. Dentists and Dental assistants need adequate information about children's oral hygiene to teach them and their parents. Also tooth brushing twice daily under parent's supervision is suggested [22].

The oldest toothbrushing method was described in 1913 by Fones and is suggested mainly for youngsters. The Bass technique places emphasis on the removal of plaque from above and slightly below the gingival margin. Bass had been changed to the Modified Bass where the bristle position and predominantly horizontal brush movements within the Bass method are retained, but vertical and sweeping motions to make circles are added. The Stillman technique is analogous to the Bass technique. The vertical motions of the Stillman technique could also be combined with the Bass, as prescribed for the Modified Bass. Charters suggested angling the comb head at 45° coronally to the margin instead of apically. Vibratory and slight rotary movement is then applied before moving to the subsequent group of teeth. An abnormal frenum may be an additive factor to plaque accumulation and may cause inhibition to proper tooth brushing. The Scrub technique is the most simple technique, with the toothbrush held parallel to the gingiva and horizontal motions to scrub the gingival crevice in an ordered fashion. There are some modification techniques such as Hirschfeld's technique which is a modification of the Fone's technique where the circular motion is smaller and concentrated over the gingival crevice. Frequency and duration of brushing are usually included with recommendations concerning the tactic of tooth brushing for children [22].

Modified Bass technique emerged as the most commonly recommended brushing method for patients aged 13 to 17 years. Oral hygiene instructions should be tailored to align with a child's developmental level and motor coordination. It is important to account for differences in brushing ability, particularly among younger children [22].

Fones technique emerged as the most commonly recommended brushing method by dentists for children aged 6 to 12 years, regardless of gender. The Modified Bass technique was the next most preferred. Ensuring proper oral hygiene during the mixed dentition stage is crucial and demands ongoing reinforcement [23].

Participants who regularly used ultrasonic toothbrushes showed a noticeable reduction in oral and salivary bacterial counts compared to those in the control group. However, proper guidance and monitoring are essential for individuals using ultrasonic toothbrushes [24].




Figure 10: Comparative Brushing Simulation with To- Fro and Oscillatory Strokes

She indicated that toothbrushes used by individuals with gingivitis had higher levels of bacterial contamination compared to those used by individuals with healthy gums. The most commonly identified microorganisms were *Staphylococcus aureus* and *Streptococcus mutans*. Toothbrushes play a significant role in the transmission of microorganisms, potentially increasing the risk of infection. Therefore, it is essential for dentists to take an active role in educating patients about the proper selection, storage, hygiene and timely replacement of toothbrushes [25].

A toothbrush is a principle instrument that helps in maintaining proper hygiene and oral care. Based on the different bristle diameters the tooth brushes have been categorized as soft (0.2 mm), medium (0.3 mm) and hard (0.4mm). Choosing the right toothbrush plays an important role in maintaining oral hygiene. Apart from choosing the right toothbrush, proper usage of the toothbrush should also be taken into consideration, as improper brushing may lead to the soft and hard tissues of the teeth. This may lead to conditions such like abrasion. Abrasion is the process in which the enamel erodes due to the force applied on teeth, improper brushing can also be caused for abrasion. Toothbrushes with different functions have been developed for oral health management. The factors that influence the surface roughness of teeth are the brushing methods, frequency, duration of brushing, bristle diameter, shape, force of brushing direction of brushing, number of bristles per tuft and its management [26].

The Modified Bass technique emerged as the most commonly recommended brushing method for patients aged 13 to 17 years. Oral hygiene instructions should be tailored to align with a child's developmental level and motor coordination. It is important to account for differences in brushing ability, particularly among younger children [22].

Tooth brushing is a very important plaque control measure. The relationship between incomplete plaque removal, sequelae of gingivitis and periodontitis and also the occurrence of dental caries has been proven. There was a wide diversity between recommendations on tooth brushing methods. Tooth brushing is the cornerstone of dental health education to prevent caries and periodontal disease. More high quality and long term studies are required to investigate the effectiveness of brushing techniques in the prevention of gingivitis, periodontitis and caries. Patients pay poor attention to the lingual sites during their regular tooth brushing practices, this may be because these sites do not affect the aesthetics and have more difficult access. So, special attention should be given to brushing techniques in lingual sites. Certain factors may influence the effectiveness of the tooth brushing technique like the dexterity of the patient, level of comprehension of the patient after demonstrating the technique, the features of the toothbrush including filament arrangement, orientation, size, shape and flexibility. Another study performed by Jagadheeswari *et al.* concluded that Modified Bass technique is the most commonly taught brushing technique to children. This study obtained results that were similar to the study done by Khalid *et al.* which also concluded that Modified Bass technique is highly effective in plaque control. In addition to this the study also concluded that Stillman's technique has lesser effectiveness in removal of plaque deposits. Advantages of this study were that it had easy access, large availability of data, similar ethnicity and high internal validity. It was also used to identify any mistakes in the brushing techniques advised. Limitation of this study was that it had low external validity. Sample size was small and inadequate. It was a uncentered study with geographic limitation. Future scope was that it should be conducted as a multi centered study with extension in the geographic limitation. And also to attain effective brushing technique to decrease the caries incidence and periodontal problems [23].

Organisms are not only associated with oral cavity but also seen in toothbrush which includes *Streptococcus mutans*, *Staphylococcus aureus*, *Pseudomonas*, *Lactobacillus*, *Klebsiella*, *Candida* species. Toothbrushes also has a significant role in disease transmission and increase the risk of infection since they can serve as are servoir for microorganisms in healthy, oral-diseased and in immunocompromised people. Contamination is the state of retention and survival of infectious organisms that occur on animate or inanimate objects. Contaminated toothbrushes may play a role in both systemic and localized diseases. This toothbrush contamination is associated with transmission of severe health problems which includes cardiovascular diseases, respiratory disorders, gastrointestinal diseases, arthritis, bacteremia, renal problems and stroke. Toothbrushes can become contaminated from the oral cavity, environment, hands, aerosol contamination and storage containers and the bacteria which attach to the toothbrush gets accumulated and survive on toothbrushes will helps in transmitting the diseases. In 1920 Cobb reported that toothbrush is the cause of repeated infections in the oral cavity. Contaminated tooth brush acts as an environment for microbial transport, growth. Toothbrush heads between the bristle tufts is a favourable medium for the growth of microorganisms. This can be the cause of reinfection of a person with pathogenic bacteria (autoinoculation) or it can acts as a significant risk of dissemination of infection for certain patients such as immunosuppressed, cardiopathic, organ transplant recipients. Various factors such as inadequate storage, the toothbrush use without decontamination, using the same toothbrush for a longer period of time without changing with new ones, survival of microorganisms for a long time leads to autoinoculation which may results in repeated entry of prospective pathogens and infection in the oral cavity especially in children and in immunocompromised patients. So, disinfection of the toothbrushes is an essential part to prevent various diseases. This condition is specifically important for children, immunocompromised patients and those are undergoing organ transplantation or chemotherapy. Actualizing the problem of toothbrushes contamination, the choice of proper tools and methods for their disinfection and the patients' education are important issues which should bring into the focus of dentists in everyday practice, because of the need of prevention the potential influence to the oral and systemic health [25].

A toothbrush is a principle instrument that helps in maintaining proper hygiene and oral care. Based on the different bristle diameters the tooth brushes have been categorized as soft (0.2 mm), medium (0.3 mm) and hard (0.4 mm). Choosing the right toothbrush plays an important role in maintaining oral hygiene. Apart from choosing the right toothbrush, proper usage of the toothbrush should also be taken into consideration, as improper brushing may lead to the soft and hard tissues of the teeth. This may lead to conditions such like abrasion. Abrasion is the process in which the enamel erodes due to the force applied on teeth, improper brushing can also be caused for abrasion. Toothbrushes with different functions have been developed

for oral health management. The factors that influence the surface roughness of teeth are the brushing methods, frequency, duration of brushing, bristle diameter, shape, force of brushing direction of brushing, number of bristles per tuft and its management [26].

Electric toothbrushes may be less effective for blind children due to their limited tactile feedback, which makes it challenging for them to feel the pressure and identify the areas being cleaned. This can hinder effective brushing techniques and result in inadequate plaque removal. Additionally, the complexity of electric toothbrushes- often relying on visual cues and features like timers and sensors- creates difficulties for blind children in establishing consistent routines and gaining confidence in their use. Caregivers and healthcare providers play a crucial role in supporting the unique needs of blind children by developing tailored oral care solutions, including the use of manual brushes and adaptive tools such as Braille instructions [27].

Manual toothbrushes were widely used due to their affordability and availability. However, their effectiveness was highly dependent on proper brushing techniques, duration and parental supervision, which could be inconsistent in younger children. Since children often lacked the necessary motor coordination for thorough plaque removal, they may have been at a higher risk of developing caries if manual brushing was not done correctly. Electric toothbrushes, including oscillating-rotating and sonic types, were introduced as alternatives to manual toothbrushes to improve plaque removal. Several studies suggested that electric toothbrushes offered better cleaning efficiency, particularly for children who struggled with manual dexterity. Research indicated that powered toothbrushes could significantly reduce plaque and gingival inflammation compared to manual brushing, making them a preferable choice for younger users. Sonic toothbrushes, a subtype of electric toothbrushes, generated high-frequency vibrations that created fluid motion, potentially enhancing plaque removal beyond direct bristle contact. Some studies suggested that sonic toothbrushes may have provided superior plaque control compared to manual and oscillating-rotating electric toothbrushes, particularly in hard- to-reach areas. Despite the growing evidence supporting electric and sonic toothbrushes, more research was needed to determine their comparative effectiveness in young children. Factors such as compliance, ease of use, comfort and parental guidance played a crucial role in determining which type of toothbrush was most effective for children in this age group [28].

## CONCLUSION

The present study demonstrates that To and Fro brushing results in a significantly greater increase in surface roughness of the molar occlusal surfaces compared to Oscillatory brushing. The mean difference in Ra values was substantially higher in the To and Fro group, with a highly significant p-value, indicating a strong statistical difference between the two brushing techniques.



Within the limitations of the study, such as sample size and *in vitro* conditions, the findings suggest that the Oscillatory brushing technique is more favorable for preserving smoother occlusal surfaces on molar teeth. This technique may exert less mechanical stress on enamel and dentin, thereby reducing the potential for surface abrasion.

The results highlight that brushing technique plays a critical role in influencing tooth surface roughness and not all techniques exert the same impact. Mechanical wear from improper brushing can lead to increased surface roughness, which in turn may contribute to plaque retention, hypersensitivity and long-term tooth structure loss.

Based on the findings, Oscillatory brushing may be recommended as the preferred technique for molar occlusal surfaces to maintain smoother tooth surfaces and minimize the risk of abrasion. These insights can inform clinical recommendations and patient education regarding optimal brushing methods for long-term oral health.

Further research involving larger samples and clinical settings is encouraged to validate these findings and explore additional variables such as brushing force, duration and frequency.

## REFERENCES

- [1] Salam, T.A.A. *et al.* "Assessment of cervical abrasion, dentin hypersensitivity and its treatment needs using the cervical abrasion index of treatment needs probe." *Cureus*, vol. 15, no. 1, January 2023, e33471. doi:10.7759/cureus.33471.
- [2] Salam, T.A.A. *et al.* "The Cervical Abrasion Index of Treatment Needs (CAITN) procedure for population groups and individuals." *Cureus*, vol. 15, no. 3, March 2023, e36324. doi:10.7759/cureus.36324.
- [3] Ali, A.S. *et al.* "Cervical abrasion, sexual dimorphism and anthropometric tooth dimension." *Journal of Pharmacy and Bioallied Sciences*, vol. 14, 2022, pp. S378–S383.
- [4] Salam, T.A.A. *et al.* "Prevalence and clinical parameters of cervical abrasion as a function of population, age, gender and toothbrushing habits: A systematic review." *World Journal of Dentistry*, vol. 10, no. 6, 2019, pp. 470–480.
- [5] Kamra, Shilpa, *et al.* "Oral Hygiene Instructions with Plaque Disclosing Agents to Improve Self-Performed Dental Plaque Control: A Case Report." *Cureus*, vol. 16, no. 10, October 2024, e72205. doi:10.7759/cureus.72205.
- [6] Ali, A.S. *et al.* "The design, development and calibration of cervical abrasion index of treatment needs probe for measurement of cervical abrasion." *Journal of Pharmacy and Bioallied Sciences*, vol. 14, 2022, pp. S384–S389.
- [7] Ramanarayanan, V. *et al.* "Measuring dental diseases: A critical review of indices in dental practice and research." *Amrita Journal of Medicine*, vol. 16, 2020, pp. 152–158.
- [8] Addy, Martin *et al.* "Dentine hypersensitivity: New perspectives on an old problem." *Journal of Dentistry*, 23 September 2014.
- [9] Grender, Julie *et al.* "An 8-week randomized controlled trial comparing the effect of a novel oscillating-rotating toothbrush versus a manual toothbrush on plaque and gingivitis." *International Dental Journal*, vol. 70, suppl. 1, April 2020, pp. S7–S15. doi:10.1111/idj.12571.
- [10] Marschner, Felix, *et al.* "Systematic review and meta-analysis on prevalence and anamnestic risk factors for erosive tooth wear in the primary dentition." *International Journal of Paediatric Dentistry*, vol. 35, no. 2, July 2024, pp. 389–404. doi:10.1111/ipd.13250.
- [11] Joiner, A. *et al.* "Tooth cleaning and tooth wear-a review." *Journal of Engineering Tribology*, vol. 224, no. 6. doi:10.1243/13506501JET671.
- [12] Shellis, R.P. and M. Addy. "The interactions between attrition, abrasion and erosion in tooth wear." *Monographs in Oral Science*, vol. 25, 2014, pp. 32–45. doi:10.1159/000359936.
- [13] Freitas, S.S. *et al.* "Dentin hypersensitivity treatment of noncarious cervical lesions-A single-blind, split-mouth study." *Brazilian Oral Research*, vol. 29, 2015, p. 45. doi:10.1590/1807-3107BOR-2015.vol29.0045.
- [14] Umberto, R. *et al.* "Treatment of Dentine Hypersensitivity by Diode Laser: A Clinical Study." *International Journal of Dentistry*, 2012, p. 858950. doi:10.1155/2012/858950.
- [15] Liu, X.X. *et al.* "Pathogenesis, Diagnosis and Management of Dentin Hypersensitivity: An Evidence-Based Overview for Dental Practitioners." *BMC Oral Health*, vol. 20, 2020, p. 220. doi:10.1186/s12903-020-01199-z.
- [16] Michael, J.A. *et al.* "Non-carious cervical lesions on permanent anterior teeth: A new morphological classification." *Australian Dental Journal*, vol. 55, 2010, pp. 134–137. doi:10.1111/j.1834-7819.2010.01228.x.
- [17] Borcic, J. *et al.* "The prevalence of non-carious cervical lesions in permanent dentition." *Journal of Oral Rehabilitation*, vol. 31, 2004, pp. 117–123. doi:10.1046/j.0305-182x.2003.01223.x.
- [18] Davari, A. *et al.* "Dentin hypersensitivity: Etiology, diagnosis and treatment; a literature review." *Journal of Dentistry (Shiraz)*, vol. 14, 2013, pp. 136–145.
- [19] Waseem, M.A. *et al.* "How to deal with cervical tooth sensitivity: A review." *Modern Research in Dentistry*, vol. 1, 2017, pp. 1–4.
- [20] Veitz-Keenan, A. *et al.* "Treatments for hypersensitive noncarious cervical lesions: A Practitioners Engaged in Applied Research and Learning (PEARL) Network Randomized Clinical Effectiveness Study." *Journal of the American Dental Association*, vol. 144, 2013, pp. 495–506. doi:10.14219/jada.archive.2013.0152.
- [21] Raja, T *et al.* "Study of Neem Fiber Composite Toothbrush – Latest Approach for the Prevention of Oral Disease." *Journal of Orofacial Sciences*, vol. 16, 2024, pp. 146–151.
- [22] Ramamoorthy, J. *et al.* "Evaluation of brushing techniques taught by dental students in children with permanent dentition." *International Journal of Dentistry and Oral Science*, vol. 8, no. 1, January 2021, pp. 1487–1491. doi:10.19070/2377-8075-21000297.
- [23] Desai, K. and V. Ravindran. "Dentist's preference of brushing technique taught to children with mixed dentition." *International Journal of Dentistry and Oral Science*, vol. 8, no. 9, September 2021, pp. 4531–4534. doi:10.19070/2377-8075-2100092.
- [24] Begum, Akifa and N.P. Muralidharan. "Assessing the Efficacy of Sonic Toothbrush in Reducing the Plaque Pathogens in Comparison with Manual Brushing."

- [25] Madhumathi, D. and Sri Sakthi. "Evaluation of difference in bacterial contamination of toothbrushes between patients with gingivitis and patients with healthy gingiva – a pilot study." *International Journal of Pharmacy Research and Technology*, vol. 9, no. 2, July–December 2019.
- [26] Sarangadharan, Varshasree *et al.* "Evaluation of surface roughness of teeth post brushing simulation with different commercially available Ultrasoft toothbrush." *HIV Nursing*, vol. 23, no. 3, 2023, pp. 146–156.
- [27] Thanalakshme, P.S. and R. Ramesh. "Comparative evaluation of the effectiveness of manual and electric toothbrushes in blind children: A Randomised controlled trial." *Journal of Clinical and Diagnostic Research*, vol. 19, no. 2, February 2025. doi:10.7860/JCDR/2025/76529.20591.
- [28] Aarthi, K. and R. Ramesh. "Comparative evaluation of plaque removal efficiency of electric, manual and sonic toothbrushes in children aged 6 to 12 years: A randomized controlled trial." *Journal of Neonatal Surgery*, vol. 14, no. 5S, March 2025, pp. 86–93.